arXiv:1211.6909v1 [math.GT] 29 Nov 2012

A Sharp Upper Bound for Region Unknotting Number of Torus Knots[☆]

Vikash. S[a], Madeti. P[b,*]

[a]Department of Mathematics, IIT Ropar, Rupnagar- 140001, India.
[b]Room No. 207, Department of Mathematics, IIT Ropar, Rupnagar - 140001, India.

Abstract

Region crossing change for a knot or a proper link is an unknotting operation. In this paper, we provide a sharp upper bound on the region unknotting number for a large class of torus knots and proper links. Also, we discuss conditions on torus links to be proper.

Keywords: Torus Knots, Region Unknotting Number, Linking Number
2000 MSC: 57M25

1. Introduction

The unknotting number $u(D)$ of a knot diagram D is the minimal number of crossing changes required to unknot D and the unknotting number $u(K)$ of a knot K is given by $u(K) = inf\{u(D) : D\ is\ a\ knot\ diagram\ of\ K\}$. It is not easy to calculate $u(K)$, because it is taken over an infinite number of diagrams of K. Unknotting number lead to consider different types of unknotting operations i.e., the local transformations on link diagrams which can transform the knot diagram into a trivial knot diagram. It was shown that $\sharp$-operation [4], δ-operation [7], 3-gon operation [9], H(n)-operation [11] and n-gon [10] operations are unknotting operations.

Recently, Ayaka Shimizu [2] proposed an unknotting operation for knots called *region crossing change*. In this operation, K to be a knot and D be any diagram of K, then a region crossing change at a region R of diagram D is defined to be the crossing changes at all the crossing points on ∂R. The region unknotting number $u_R(D)$ of a knot diagram D is the minimal number of region crossing changes required to transform D into a diagram of the trivial knot without Reidemeister moves. The region unknotting number $u_R(K)$ of K is defined to be the minimal $u_R(D)$ taken over all minimal diagrams D of K.

In general, for links, the region crossing change is not an unknotting operation. In [2], Ayaka Shimizu considered the standrard diagram of Hopf link and showed that it cannot be transformed into a diagram of trivial link by region crossing changes. In [8] and [12] Cheng Zhiyun defined a link $L = K_1 \bigcup K_2 \bigcup \cdots \bigcup K_n$ to be proper if for each i, $1 \leq i \leq n$, $\sum_{j \neq i} lk(K_i, K_j) \equiv 0\ (mod\ 2)$ and proved that *region crossing change on a link is an unknotting operation if and only if the link is proper.*

The region unknotting number is intuitive and attractive for knot theorists and it is not surprising that there is no algorithm to compute it for a given knot or proper link at the moment. In [2], [3], Ayaka Shimizu showed that for a twist knot K, $u_R(K) = 1$ and for torus knots of type

---

[☆] This document is a collaborative effort.

[*] Corresponding author

Email address: `prabhakar@iitrpr.ac.in` (Madeti. P)
URL: `http://www.iitrpr.ac.in/html/faculty/prabhakar.shtml` (Madeti. P)

$K(2, 4m \pm 1)$, $u_R(K(2, 4m \pm 1)) = m$, where $m \in \mathbb{Z}^+$. As the unknotting number for torus knots is already known [5], [6], it would be worth attempting to find the region unknotting number for torus knots. In this paper, we discuss the condition for torus links to be proper and provide a sharp upper bound for a large class of torus knots and proper torus links, including all $2-, 3-, 4-, 5-, 6-$ and $7-$braid torus knots/links.

In Section 2, we discuss the condition for torus links to be proper. In Section 3, we discuss basic results, depending on elementary braid relations, which helps to provide a sharp upper bound for region unknotting number for torus links. In Section 4, we provide a sharp upper bound for region unknotting number of $K(p, np + a)$ torus knots/links, where $a = 1, 2, 3, 4, -1, -2$, and partially for $K(p, np + 5)$ type of torus links.

2. Conditions for torus link to be proper

In [12], Cheng Zhiyun showed that, *region crossing change is an unknotting operation on a link* $L = K_1 \bigcup K_2 \bigcup \cdots \bigcup K_n$ *if and only if* $\sum_{j \neq i} lk(K_i, K_j) \equiv 0 \ (mod\ 2)$ *for each* $1 \leq i \leq n$. If $gcd(p, q) = d$, we use the notation $K(p, q) = K_1 \bigcup K_2 \bigcup \cdots \bigcup K_d$, where each K_i is a component of $K(p, q)$. To provide the condition for torus links to be proper, we consider the diagram for $K(p, q)$ as a closure of toric braid $(\sigma_1 \sigma_2 \cdots \sigma_{p-1})^q$. Also, we denote $\sum_{i=1}^{n} i$ by $\sum n$. Given a d-component torus link $K(p, q)$, the number of crossings between any two components is same i.e., for $i \neq j$, $l \neq m$, $\sharp(K_i \bigcap K_j) = \sharp(K_l \bigcap K_m)$. Also, it is worth noting that, for $i \neq j$, $lk(K_i, K_j) = \frac{1}{2} \sharp(K_i \bigcap K_j)$.

Theorem 2.1. *A torus link* $K(p, q)$ *is proper iff*

$$\frac{pq(d-1)}{d^2} \equiv 0 (mod\ 2),$$

where: d=gcd(p,q).

Proof. Consider a d-component torus link $K(p, q)$. Since we are considering $K(p, q)$ as a closure of toric braid $(\sigma_1 \sigma_2 \cdots \sigma_{p-1})^q$, each component K_i contains p/d strands. For each bracket $(\sigma_1 \sigma_2 \cdots \sigma_{p-1})$, there exists only one i such that $\sharp(K_i \bigcap K_i) = (\frac{p}{d} - 1)$. Hence the total number of self-intersections is

$$\sum_{i=1}^{d} \sharp \left(K_i \bigcap K_i \right) = q \left(\frac{p}{d} - 1 \right).$$

Since,

$$\sum_{i \neq j} \sharp(K_i \bigcap K_j) = total\ no\ of\ crossings - \sum_{i=1}^{d} \sharp(K_i \bigcap K_i),$$

we have

$$\sum_{i \neq j} \sharp(K_i \bigcap K_j) = \frac{pq(d-1)}{d}. \tag{1}$$

Because $\sharp(K_i \bigcap K_j) = \sharp(K_l \bigcap K_m)$, for any $i \neq j, l \neq m$, we have

$$\sum_{i \neq j} \sharp(K_i \bigcap K_j) = \frac{(d-1)d}{2} \left(\sharp(K_1 \bigcap K_2) \right). \tag{2}$$

Thus, from (1) and (2), for any $i \neq j$

$$\sharp(K_i \bigcap K_j) = \frac{2pq}{d^2}.$$

Since all crossings in $K(p,q)$ are positive, for fixed n

$$\begin{aligned}\sum_{j\neq n} lk(K_n, K_j) &= \frac{1}{2}\sum_{j\neq n}\sharp(K_n \bigcap K_j)\\ &= \frac{1}{2}(d-1)\left(\sharp(K_i \bigcap K_j)\right)\\ &= \frac{pq(d-1)}{d^2}.\end{aligned}$$

Hence, a torus link $K(p,q)$, where d=gcd(p,q), is proper iff

$$\frac{pq(d-1)}{d^2} \equiv 0(mod\ 2). \qquad \square$$

Corollary 2.1. *Let $K(p,q)$ be any d-componant torus link where $p = 2^m k$ and $q = 2^n k'$, where m and n are the highest powers of 2 in p and q, respectively. Then $K(p,q)$ is proper iff either $m = n = 0$ or $m \neq n$.*

3. Preliminaries

To find the sharp upper bounds on the region unknotting number of torus links, we use the following:

Theorem 3.1. *([1], Theorem 3.1) For every p, the p-braid*

$$\underbrace{\sigma_1\sigma_2\cdots\sigma_{p-1}}_{1}\underbrace{\sigma_1\sigma_2\cdots\sigma_{p-2}\sigma_{p-1}^{-1}}_{2}\underbrace{\sigma_1\sigma_2\cdots\sigma_{p-2}^{-1}\sigma_{p-1}^{-1}}_{3}\cdots\underbrace{\sigma_1^{-1}\sigma_2^{-2}\cdots\sigma_{p-1}^{-1}}_{p}$$

is a trivial p-braid.

Remark 1. Using Theorem 3.1, we observe the following:
(i) For every p, the p-braid

$$\underbrace{\sigma_1^{-1}\sigma_2^{-2}\cdots\sigma_{p-1}^{-1}}_{1}\underbrace{\sigma_1^{-1}\sigma_2^{-2}\cdots\sigma_{p-2}^{-1}\sigma_{p-1}}_{2}\cdots\underbrace{\sigma_1^{-1}\sigma_2\cdots\sigma_{p-1}}_{p-1}\underbrace{\sigma_1\sigma_2\cdots\sigma_{p-1}}_{p}$$

is a trivial p-braid.
(ii) The closure of $n+1$ braids

$$\underbrace{\sigma_1\sigma_2\cdots\sigma_{n-1}\sigma_n^{-1}}_{1}\underbrace{\sigma_1\sigma_2\cdots\sigma_{n-1}^{-1}\sigma_n^{-1}}_{2}\cdots\underbrace{\sigma_1^{-1}\sigma_2^{-1}\cdots\sigma_n^{-1}}_{n}$$

and

$$\underbrace{\sigma_1\sigma_2\cdots\sigma_{n-1}\sigma_n}_{1}\underbrace{\sigma_1\sigma_2\cdots\sigma_{n-1}\sigma_n^{-1}}_{2}\cdots\underbrace{\sigma_1\sigma_2^{-1}\cdots\sigma_n^{-1}}_{n}$$

are trivial links.

Lemma 3.1. *For every p and a, where $p > a$, the p-braid*
$\underbrace{\eta_1\kappa_{p-1}}_{1}\underbrace{\eta_2\kappa_{p-2}\sigma_{p-1}^{-1}}_{2}\cdots\underbrace{\eta_{a-1}\kappa_{p-a+1}\sigma_{p-a+2}^{-1}\cdots\sigma_{p-1}^{-1}}_{a-1}\underbrace{\eta_a\sigma_{p-a+1}^{-1}\cdots\sigma_{p-1}^{-1}}_{a}$ *is Markov equivalent of* $\eta'_1\eta'_2\cdots\eta'_a$, *where* $\eta'_i = \sigma_1^{g_{i,1}}\sigma_2^{g_{i,2}}\cdots\sigma_{p-a-1}^{g_{i,p-a-1}}$, $\eta_i = \eta'_i\sigma_{p-a}^{g_{i,p-a}}$ *with* $g_{i,j} = \pm 1$, *for* $i = 1, 2, \ldots, a$; $j = 1, 2, \ldots, p-a$ *and* $\kappa_j = \sigma_{p-a+1}\sigma_{p-a+2}\cdots\sigma_j$.

Proof. Using fundamental braid relations, we observe that

$$\begin{aligned}
&\underbrace{\eta_1\kappa_{p-1}}_{1}\underbrace{\eta_2\kappa_{p-2}\sigma_{p-1}^{-1}}_{2}\cdots\underbrace{\eta_{a-1}\kappa_{p-a+1}\sigma_{p-a+2}^{-1}\cdots\sigma_{p-1}^{-1}}_{a-1}\underbrace{\eta_a\sigma_{p-a+1}^{-1}\cdots\sigma_{p-1}^{-1}}_{a}\\
&\sim_M \underbrace{\eta'_1\sigma_{p-a}^{g_{1,p-a}}\kappa_{p-1}}_{1}\underbrace{\eta'_2\sigma_{p-a}^{g_{2,p-a}}\kappa_{p-2}\sigma_{p-1}^{-1}}_{2}\cdots\underbrace{\eta'_a\sigma_{p-a}^{g_{a,p-a}}\sigma_{p-a+1}^{-1}\cdots\sigma_{p-1}^{-1}}_{a}\\
&\sim_M \underbrace{\eta'_1\sigma_{p-a}^{g_{1,p-a}}\kappa_{p-2}}_{1}\underbrace{\eta'_2\sigma_{p-a}^{g_{2,p-a}}\kappa_{p-3}\sigma_{p-2}^{-1}}_{2}\underline{\sigma_{p-1}\sigma_{p-2}}\cdots\underbrace{\eta'_a\sigma_{p-a}^{g_{a,p-a}}\sigma_{p-a+1}^{-1}\cdots\sigma_{p-1}^{-1}}_{a}\\
&\qquad\vdots\\
&\sim_M \underbrace{\eta'_1\sigma_{p-a}^{g_{1,p-a}}\kappa_{p-2}}_{1}\underbrace{\eta'_2\sigma_{p-a}^{g_{2,p-a}}\kappa_{p-3}\sigma_{p-2}^{-1}}_{2}\cdots\underbrace{\eta'_a\sigma_{p-a}^{-1}\cdots\sigma_{p-2}^{-1}}_{a}\underline{\sigma_{p-1}^{g_{a,p-a}}\sigma_{p-2}\cdots\sigma_{p-a}}\\
&\sim_M \underbrace{\eta'_1\sigma_{p-a}^{g_{1,p-a}}\kappa_{p-2}}_{1}\underbrace{\eta'_2\sigma_{p-a}^{g_{2,p-a}}\kappa_{p-3}\sigma_{p-2}^{-1}}_{2}\cdots\underbrace{\eta'_{a-1}\sigma_{p-a}^{g_{a-1,p-a}}\sigma_{p-a+1}^{-1}\sigma_{p-a+2}^{-1}\cdots\sigma_{p-2}^{-1}}_{a-1}\eta'_a.
\end{aligned}$$

Continuing the same procedure after $a-2$ steps, we obtain

$$\sim_M \underbrace{\eta'_1\sigma_{p-a}^{g_{1,p-a}}\kappa_{p-3}}_{1}\underbrace{\eta'_2\sigma_{p-a}^{g_{2,p-a}}\kappa_{p-4}\sigma_{p-3}^{-1}}_{2}\cdots\underbrace{\eta'_{a-2}\sigma_{p-a}^{g_{a-2,p-a}}\sigma_{p-a+1}^{-1}\cdots\sigma_{p-3}^{-1}}_{a-2}\eta'_{a-1}\eta'_a.$$

After $\sum(a-3)$ steps, this p-braid is Markov equivalent of $\eta'_1\eta'_2\cdots\eta'_a$. □

Lemma 3.2. *For any two even numbers p and q with $p > q$,*

$$\eta_1\kappa_1\eta_2\kappa_2\cdots\eta_q\kappa_q \sim_M \eta'_1\eta'_2\cdots\eta'_q,$$

where $\eta_i = \eta'_i\sigma_{p-q}^{g_{i,p-q}}$ *and* $\eta'_i = \sigma_1^{g_{i,1}}\sigma_2^{g_{i,2}}\cdots\sigma_{p-q-1}^{g_{i,p-q-1}}$; *and* $\kappa_i = \sigma_{p-q+1}^{-1}\sigma_{p-q+2}^{-1}\cdots\sigma_{p-i}^{-1}\sigma_{p-i+1}\cdots\sigma_{p-1}$ *with* $g_{i,j} = \pm 1$ *for* $i = 1, 2, \cdots, q$ *and* $j = 1, 2, \cdots, p-q$.

Proof. Using fundamental braid relations we observe that

$$\begin{aligned}
&\underbrace{\eta_1\sigma_{p-q+1}^{-1}\cdots\sigma_{p-1}^{-1}}_{1}\underbrace{\eta_2\sigma_{p-q+1}^{-1}\cdots\sigma_{p-2}^{-1}\sigma_{p-1}}_{2}\cdots\underbrace{\eta_q\sigma_{p-q+1}\cdots\sigma_{p-1}}_{q}\\
&\sim_M \underbrace{\eta_1\sigma_{p-q+1}^{-1}\cdots\sigma_{p-2}^{-1}}_{1}\underbrace{\eta_2\sigma_{p-q+1}^{-1}\cdots\sigma_{p-3}^{-1}\sigma_{p-2}}_{2}\underline{\sigma_{p-1}^{-1}\sigma_{p-2}^{-1}}\cdots\underbrace{\eta_q\sigma_{p-q+1}\cdots\sigma_{p-1}}_{q}\\
&\qquad\vdots\\
&\sim_M \underbrace{\eta_1\sigma_{p-q+1}^{-1}\cdots\sigma_{p-2}^{-1}}_{1}\underbrace{\eta_2\sigma_{p-q+1}^{-1}\cdots\sigma_{p-3}^{-1}\sigma_{p-2}}_{2}\cdots\underbrace{\eta'_q\sigma_{p-q}\cdots\sigma_{p-2}}_{q}\underline{\sigma_{p-1}^{-1}\sigma_{p-2}^{-1}\cdots\sigma_{p-q}^{-1}}\\
&\sim_M \underbrace{\eta_1\sigma_{p-q+1}^{-1}\cdots\sigma_{p-2}^{-1}}_{1}\underbrace{\eta_2\sigma_{p-q+1}^{-1}\cdots\sigma_{p-3}^{-1}\sigma_{p-2}}_{2}\cdots\underbrace{\eta_{q-1}\sigma_{p-q+1}\cdots\sigma_{p-2}}_{q-1}\eta'_q.
\end{aligned}$$

Continuing the same procedure after $q-2$ steps, we obtain

$$\sim_M \underbrace{\eta_1\sigma_{p-q+1}^{-1}\cdots\sigma_{p-3}^{-1}}_{1}\underbrace{\eta_2\sigma_{p-q+1}^{-1}\cdots\sigma_{p-4}^{-1}\sigma_{p-3}}_{2}\cdots\underbrace{\eta_{q-2}\sigma_{p-q+1}\cdots\sigma_{p-3}}_{q-1}\eta'_{q-1}\eta_q.$$

After $\sum(q-3)$ similar steps, this p-braid is Markov equivalent of $\eta'_1\eta'_2\cdots\eta'_q$. $\square$

Lemma 3.3. *Let* $\beta_j=\sigma_1^{g_{j,1}}\sigma_2^{g_{j,2}}\cdots\sigma_{p-5}^{g_{j,p-5}}$, *where* $g_{j,k}=\pm1$, $1\le j\le 2$. *Then*

$$\beta_1\sigma_{p-4}^{g_{1,p-4}}\sigma_{p-3}\sigma_{p-2}\sigma_{p-1}^{-1}\beta_2\sigma_{p-4}^{g_{2,p-4}}\sigma_{p-3}\sigma_{p-2}^{-1}\sigma_{p-1}^{-1}\sim_M\beta_1\beta_2.$$

Proof. Consider that

$$\begin{aligned}
L.H.S &\sim_M \beta_1\sigma_{p-4}^{g_{1,p-4}}\sigma_{p-3}\sigma_{p-2}\beta_2\sigma_{p-4}^{g_{2,p-4}}\sigma_{p-3}\sigma_{p-2}^{-1}\underline{\sigma_{p-1}^{-1}\sigma_{p-2}^{-1}}\\
&\sim_M \beta_1\sigma_{p-4}^{g_{1,p-4}}\sigma_{p-3}\beta_2\sigma_{p-4}^{g_{2,p-4}}\sigma_{p-3}^{-1}\underline{\sigma_{p-3}^{-1}\sigma_{p-2}\sigma_{p-3}}\\
&\sim_M \beta_1\sigma_{p-4}^{g_{1,p-4}}\beta_2\underline{\sigma_{p-4}^{-1}\sigma_{p-3}^{g_{2,p-4}}\sigma_{p-4}}\\
&\sim_M \beta_1\beta_2.
\end{aligned}$$

Lemma 3.4. *If* $p(\ge 4)\equiv 0$ *or* ±2 $(mod\ 6)$, *then the closure of* $\sigma_1^{-1}\sigma_2^{-1}\cdots\sigma_{p-1}^{-1}\sigma_1^{g_{2,1}}\sigma_2^{g_{2,2}}\cdots\sigma_{p-1}^{g_{2,p-1}}$ $\sigma_1^{g_{3,1}}\sigma_2^{g_{3,2}}\cdots\sigma_{p-1}^{g_{3,p-1}}$, *is a trivial link. Here*
$g_{2,j}=-1$ *for* $j\in\{p-2,p-3,p-8,p-9,\cdots,p-2-6(\lfloor\frac{p+2}{6}\rfloor-1),p-3-6(\lfloor\frac{p-4}{6}\rfloor)\}$,
$g_{3,j}=-1$ *for* $j\in\{p-3,p-4,p-9,p-10,\cdots,p-3-6(\lfloor\frac{p+2}{6}\rfloor-1),p-4-6(\lfloor\frac{p-4}{6}\rfloor)\}$.

Proof. By using fundamental braid relations, we obeserve that the p-braid $\sigma_1^{-1}\sigma_2^{-1}\cdots\sigma_{p-1}^{-1}\sigma_1^{g_{2,1}}$ $\sigma_2^{g_{2,2}}\cdots\sigma_{p-1}^{g_{2,p-1}}\sigma_1^{g_{3,1}}\sigma_2^{g_{3,2}}\cdots\sigma_{p-1}^{g_{3,p-1}}$ is Markov equivalent of
Case 1. If $p\equiv 0$ $(mod\ 6)$,

$$\begin{aligned}
&\underbrace{\sigma_1^{-1}\sigma_2^{-1}\cdots\sigma_{p-2}^{-1}}_{1}\underbrace{\sigma_1\sigma_2\cdots\sigma_{p-3}^{-1}\sigma_{p-2}}_{2}\underbrace{\sigma_1\sigma_2^{-1}\cdots\sigma_{p-3}\sigma_{p-2}}_{3}\underline{\sigma_{p-1}^{-1}\sigma_{p-2}^{-1}\sigma_{p-3}^{-1}}\\
&\sim_M \underbrace{\sigma_1^{-1}\sigma_2^{-1}\cdots\sigma_{p-3}^{-1}}_{1}\underbrace{\sigma_1\sigma_2\sigma_3^{-1}\cdots\sigma_{p-4}\sigma_{p-3}}_{2}\underline{\sigma_{p-2}^{-1}\sigma_{p-3}^{-1}}\underbrace{\sigma_1\sigma_2^{-1}\cdots\sigma_{p-4}^{-1}}_{3}\\
&\sim_M \underbrace{\sigma_1^{-1}\sigma_2^{-1}\cdots\sigma_{p-4}^{-1}}_{1}\underbrace{\sigma_1\sigma_2\cdots\sigma_{p-4}}_{2}\underbrace{\sigma_1\sigma_2^{-1}\cdots\sigma_{p-9}^{-1}\sigma_{p-8}\cdots\sigma_{p-5}\sigma_{p-4}^{-1}}_{3}\\
&\sim_M \underbrace{\sigma_1^{-1}\sigma_2^{-1}\cdots\sigma_{p-5}^{-1}}_{1}\underbrace{\sigma_1\sigma_2\sigma_3^{-1}\cdots\sigma_{p-5}}_{2}\underbrace{\sigma_1\sigma_2^{-1}\cdots\sigma_{p-7}\sigma_{p-6}\sigma_{p-5}^{-1}}_{3}\underline{\sigma_{p-4}\sigma_{p-5}\sigma_{p-6}^{-1}}\\
&\sim_M \underbrace{\sigma_1^{-1}\sigma_2^{-1}\cdots\sigma_{p-6}^{-1}}_{1}\underbrace{\sigma_1\sigma_2\sigma_3^{-1}\cdots\sigma_{p-7}\sigma_{p-6}}_{2}\underline{\sigma_{p-5}\sigma_{p-6}^{-1}}\underbrace{\sigma_1\sigma_2^{-1}\cdots\sigma_{p-8}\sigma_{p-7}}_{3}\\
&\sim_M \underbrace{\sigma_1^{-1}\sigma_2^{-1}\cdots\sigma_{p-7}^{-1}}_{1}\underbrace{\sigma_1\sigma_2\cdots\sigma_{p-9}^{-1}\sigma_{p-8}^{-1}\sigma_{p-7}}_{2}\underbrace{\sigma_1\sigma_2^{-1}\cdots\sigma_{p-10}^{-1}\sigma_{p-9}^{-1}\sigma_{p-8}\sigma_{p-7}}_{3}\\
&\qquad\vdots\\
&\sim_M \sigma_1^{-1}\sigma_2^{-1}\sigma_3^{-1}\sigma_4^{-1}\sigma_5^{-1}\sigma_1\sigma_2\sigma_3^{-1}\sigma_4^{-1}\sigma_5\sigma_1\sigma_2^{-1}\sigma_3^{-1}\sigma_4\sigma_5.
\end{aligned}$$

Case 2. If $p\equiv 2$ $(mod\ 6)$, then

$$\sigma_1^{-1}\sigma_2^{-1}\sigma_3^{-1}\sigma_4^{-1}\sigma_5^{-1}\sigma_6^{-1}\sigma_7^{-1}\sigma_1\sigma_2\sigma_3\sigma_4\sigma_5^{-1}\sigma_6^{-1}\sigma_7\sigma_1\sigma_2\sigma_3\sigma_4^{-1}\sigma_5^{-1}\sigma_6\sigma_7.$$

Case 3. If $p \equiv -2 \ (mod\ 6)$, then

$$\sigma_1^{-1}\sigma_2^{-1}\sigma_3^{-1}\sigma_1^{-1}\sigma_2^{-1}\sigma_3\sigma_1^{-1}\sigma_2\sigma_3.$$

It is easy to observe that closures of these braids are trivial links. □

Lemma 3.5. *The closure of the braids*

$$\sigma_i\sigma_{i+1}\cdots\sigma_j\sigma_i^{-1}\sigma_{i+1}^{-1}\cdots\sigma_j^{-1}; if\ \ i \leq j$$

and

$$\sigma_i\sigma_{i-1}\cdots\sigma_j\sigma_i^{-1}\sigma_{i-1}^{-1}\cdots\sigma_j^{-1}; if\ \ i \geq j$$

are trivial links.

Proof. The proof directly follows from the elementary braid relations.

Remark 2. If we take $i = 1$ and $j = n$ in Lemma 3.5, then the closures of $n+1$-braids

$$\sigma_1\sigma_2\cdots\sigma_n\sigma_1^{-1}\sigma_2^{-1}\cdots\sigma_n^{-1}$$

and

$$\sigma_1^{-1}\sigma_2^{-1}\cdots\sigma_n^{-1}\sigma_1\sigma_2\cdots\sigma_n$$

are trivial links.

4. Sharp upper bound for region unknotting number of torus knots

Let K be any knot/link and $c(K)$ be the crossings number of k. Then by [2], the trivial upper bound for region unknotting number of knot/link K is $u_R(K) = (c(K)+2)/2$. In case of torus knots/links $K(p,q)$, this trivial upper bound is $u_R(K) = ((p-1)q+2)/2$.

In most cases, to find a sharp upper bound for region unknotting number of torus links $K(p,q)$ with $p < q$, we consider $K(p,q)$ as a closure of $(\sigma_1\sigma_2\cdots\sigma_{p-1})^q$. In particular, we consider the diagram and mark the regions as shown in Figure 1(a). Another diagram we consider for $K(p,q)$ where (q=np+4), is the closure of the braid $(\sigma_1\sigma_2\cdots\sigma_{p-1})^{np}(\sigma_1\sigma_2\cdots\sigma_3)^a \underbrace{\sigma_4\sigma_3\cdots\sigma_1}\underbrace{\sigma_5\sigma_4\cdots\sigma_2}\cdots \cdots\underbrace{\sigma_{p-1}\sigma_{p-2}\cdots\sigma_{p-4}}$ as shown in Figure 1(b), where c and c' are crossing numbers of $K(p,np+4)$ and $K(p,np)$, respectively.

Let $X_i = \{2i(p-1), 2i(p-1)-2, \cdots, 2i(p-1)-2(i-1)\}$ be the set of i regions in $(\sigma_1\sigma_2\cdots\sigma_{p-1})^q$. We use $X_i^{'s}$ to identify the selected regions throughout the paper.

Theorem 4.1. *For a torus link $K(p,q)$, where $q = np$ or $np+1$, we obtain the following:*

1. *If p is odd, then*
$$u_R(K(p,q)) \leq \frac{n(p^2-1)}{8}.$$
2. *If p is even and n is odd, then $K(p,np)$ is not proper and*
$$u_R(K(p,np+1)) \leq \frac{np^2+2p}{8}.$$



Figure 1: Minimal diagrams for (a) $K(7,7)$; (b) $K(7,7n+4)$

3. *If both p and n are even, then*

$$u_R(K(p,q)) \leq \frac{np^2}{8}.$$

PROOF. Consider $K(p,q)$ as the closure of $(\sigma_1\sigma_2\cdots\sigma_{p-1})^q$, as shown in Figure 1(a). Then,
Case 1. If p is odd, then we select $\{(i-1)(p-1)p+\{X_1,X_2,\cdots,X_{\frac{p-1}{2}}\}\}_{i=1}^n$ regions so that after making region crossing change on these regions, we obtain

$$(\mu_1\mu_2\cdots\mu_p)^n; if q = np,$$
$$(\mu_1\mu_2\cdots\mu_p)^n\mu_1; if q = np+1$$

where $\mu_i = \sigma_1\sigma_2\cdots\sigma_{p-i}\sigma_{p-i+1}^{-1}\sigma_{p-i+2}^{-1}\cdots\sigma_{p-1}^{-1}$. But $(\mu_1\mu_2\cdots\mu_p)^n$ is a trivial p-braid by Theorem 3.1. Hence, the closure of the resultant braid is a trivial link.

Observe that the number of region crossing changes is equal to $n\sum(\frac{p-1}{2}) = \frac{n(p^2-1)}{8}$. So

$$u_R(K(p,np)) \leq \frac{n(p^2-1)}{8}.$$

Case 2. If p is even and n is odd, then that $K(p,np)$ is not proper follows from Theorem 2.1 and for $K(p,np+1)$ we select the regions $\{(i-1)(p-1)2p+\{X_1,X_2,\cdots,X_{\frac{p}{2}},((p-1)(2p-1)+1)-\{X_1,X_2,\cdots,X_{\frac{p-2}{2}}\}\}\}_{i=1}^{\frac{n-1}{2}},(n-1)p(p-1)+\{X_1,X_2,\cdots,X_{\frac{p}{2}}\}$. Then, after making region crossing change on these regions, we obtain

$$(\mu_1\mu_2\cdots\mu_p\nu_1\nu_2\cdots\nu_p)^{\frac{n-1}{2}}\mu_1\mu_2\cdots\mu_{p-1}\mu_p\mu_p$$

where $\mu_i = \sigma_1\sigma_2\cdots\sigma_{p-i}\sigma_{p-i+1}^{-1}\sigma_{p-i+2}^{-1}\cdots\sigma_{p-1}^{-1}$ and $\nu_i = \sigma_1^{-1}\sigma_2^{-1}\cdots\sigma_{p-i}^{-1}\sigma_{p-i+1}\cdots\sigma_{p-1}$. Note that ν_i is a mirror image of μ_i. Observe that $\mu_1\mu_2\cdots\mu_p$ and $\nu_1\nu_2\cdots\nu_p$ are trivial p-braids by Theorem 3.1 and Remark 1. Hence, the closure of the resultant braid is a trivial link.



Figure 2: Minimal diagrams for $K(2, q)$

Observe that the number of region crossing changes is equal to $\frac{(n-1)}{2}(\sum \frac{p}{2} + \sum \frac{p-2}{2}) + \sum \frac{p}{2} = \frac{np^2+2p}{8}$. Thus,

$$u_R(K(p, np+1)) \leq \frac{np^2+2p}{8}.$$

Case 3. If both p and n are even, then we select $\{(i-1)(p-1)2p + \{X_1, X_2, \cdots, X_{\frac{p}{2}}, ((p-1)(2p-1)+1) - \{X_1, X_2, \cdots, X_{\frac{p-2}{2}}\}\}\}_{i=1}^{\frac{n}{2}}$ regions so that after making region crossing change on these regions, we obtain

$$(\mu_1\mu_2\cdots\mu_p\nu_1\nu_2\cdots\nu_p)^{\frac{n}{2}}; if q = np,$$
$$(\mu_1\mu_2\cdots\mu_p\nu_1\nu_2\cdots\nu_p)^{\frac{n}{2}}\mu_1; if q = np+1$$

where $\mu_i = \sigma_1\sigma_2\cdots\sigma_{p-i}\sigma_{p-i+1}^{-1}\sigma_{p-i+2}^{-1}\cdots\sigma_{p-1}^{-1}$ and $\nu_i = \sigma_1^{-1}\sigma_2^{-1}\cdots\sigma_{p-i}^{-1}\sigma_{p-i+1}\cdots\sigma_{p-1}$. So by Theorem 3.1 and Remark 1, the closure of the resultant braid is a trivial link.

Observe that the number of region crossing changes is equal to $\frac{n}{2}(\sum \frac{p}{2} + \sum \frac{p-2}{2}) = \frac{np^2}{8}$. Thus

$$u_R(K(p, np)) \leq \frac{np^2}{8}. \qquad \square$$

Observe that the unknotting number of $K(p, np)$ and $K(p, np+1)$ are identical. Also, the number of crossing changes due to the region crossing changes prescribed in Theorem 4.1 for proper $K(p, np)$ and $K(p, np+1)$ torus links is equal to the unknotting number for $K(p, np)$.

Remark 3. Consider a proper $K(2, q)$ torus knot/link then, the only possible minimal diagrams for $K(2, q)$ are the closures of σ_1^q, as shown in Figure 2. In any case, the region crossing change at $(q+1)^{th}$ or $(q+2)^{th}$ region provides us the mirror image of $K(2, q)$. The region crossing change at any other region provides us $K(2, q-4)$. Hence, it is easy to observe that $u_R(K(2, q)) = \lfloor \frac{q+2}{4} \rfloor$.

Theorem 4.2. *For torus links $K(p, np+a)$, where a is odd and $p \equiv 0$ or $\pm 1 \pmod{a}$, $n \geq 1$. Then, we obtain the following:*

1. *If p is odd, then*

$$u_R(K(p, np+a)) \leq \frac{n(p^2-1)}{8} + \left\lfloor \frac{p+1}{a} \right\rfloor \left(\frac{a^2-1}{8} \right).$$

2. *If both p and n are even, then*

$$u_R(K(p, np+a)) \leq \frac{np^2}{8} + \left\lfloor \frac{p+1}{a} \right\rfloor \left(\frac{a^2-1}{8} \right).$$

Proof. Consider $K(p, np+a)$ as the closure of $(\sigma_1\sigma_2\cdots\sigma_{p-1})^{np+a}$ as in Figure 1(a). Then,

Case 1. If p is odd, then we select $\{\{(i-1)(p-1)p+\{X_1, X_2, \cdots, X_{\frac{p-1}{2}}\}\}_{i=1}^{n}, \{(np(p-1)-ja)+\{X_1, X_2, \cdots, X_{\frac{a-1}{2}}\}\}_{j=0}^{\lfloor\frac{p+1}{a}\rfloor-1}\}$ regions so that after making region crossing changes at these regions, the closure of the resultant braid is a trivial link by Theorem 4.1, Remark 1 and Lemma 3.1. Observe that the number of the regions selected is $n\sum\frac{p-1}{2}+\lfloor\frac{p+1}{a}\rfloor\sum\frac{a-1}{2}=\frac{n(p^2-1)}{8}+\lfloor\frac{p+1}{a}\rfloor\frac{a^2-1}{8}$. Thus,

$$u_R(K(p, np+a)) \leq \frac{n(p^2-1)}{8}+\left\lfloor\frac{p+1}{a}\right\rfloor\left(\frac{a^2-1}{8}\right).$$

Case 2. If both p and n are even, then we select $\{\{(i-1)(p-1)2p+\{X_1, X_2, \cdots, X_{\frac{p}{2}}, ((p-1)(2p-1)+1)-\{X_1, X_2, \cdots, X_{\frac{p-2}{2}}\}\}\}_{i=1}^{\frac{n}{2}}, \{(np(p-1)-ja)+\{X_1, X_2, \cdots, X_{\frac{a-1}{2}}\}\}_{j=0}^{\lfloor\frac{p+1}{a}\rfloor-1}\}$ regions so that after making region crossing changes at these regions, the closure of the resultant braid is a trivial link by Theorem 4.1, Remark 1 and Lemma 3.1. Observe that the number of regions selected is $\frac{n}{2}(\sum\frac{p}{2}+\sum\frac{p-2}{2})+\lfloor\frac{p+1}{a}\rfloor\sum\frac{a-1}{2}=\frac{np^2}{8}+\lfloor\frac{p+1}{a}\rfloor\frac{a^2-1}{8}$. Thus,

$$u_R(K(p, np+a)) \leq \frac{np^2}{8}+\left\lfloor\frac{p+1}{a}\right\rfloor\left(\frac{a^2-1}{8}\right).$$

□

Theorem 4.3. *For torus links $K(p, np+a)$, where a is odd and $p \equiv 2 \pmod a$, $n \geq 1$, we obtain the following:*

1. *If p is odd, then*

$$u_R(K(p, np+a)) \leq \frac{n(p^2-1)}{8}+\frac{p-2}{a}\left(\frac{a^2-1}{8}\right)+\left\lfloor\frac{a+2}{4}\right\rfloor.$$

2. *If both p and n are even, then*

$$u_R(K(p, np+a)) \leq \frac{np^2}{8}+\frac{p-2}{a}\left(\frac{a^2-1}{8}\right)+\left\lfloor\frac{a+2}{4}\right\rfloor.$$

Proof. Consider $K(p, np+a)$ as the closure of $(\sigma_1\sigma_2\cdots\sigma_{p-1})^{np+a}$ as in Figure 1(a). Then,

Case 1. If both p and a are odd, then $\exists\ m \in \mathbb{N}\cup\{0\}$ such that $a = 4m \pm 1$. We select $\{\{(i-1)(p-1)p+\{X_1, X_2, \cdots, X_{\frac{p-1}{2}}\}\}_{i=1}^{n}, \{(np(p-1)-ja)+\{X_1, X_2, \cdots, X_{\frac{a-1}{2}}\}\}_{j=0}^{\frac{p-2}{a}-1}, \{(np+1)(p-1)+1+4k(p-1)\}_{k=0}^{m-1}\}$ regions so that after making region crossing changes at these regions, the closure of the resultant braid is a trivial link by Theorem 4.1 and Lemma 3.1. Observe that the number of selected regions is $n\sum\frac{p-1}{2}+\frac{p-2}{a}\sum\frac{a-1}{2}+m=n\frac{p^2-1}{8}+\frac{p-2}{a}(\frac{a^2-1}{8})+\lfloor\frac{a+2}{4}\rfloor$. Thus,

$$u_R(K(p, np+a)) \leq \frac{n(p^2-1)}{8}+\frac{p-2}{a}\left(\frac{a^2-1}{8}\right)+\left\lfloor\frac{a+2}{4}\right\rfloor.$$

Case 2. If both p and n are even and a is odd, then $\exists\ m \in \mathbb{N}\cup\{0\}$ such that $a = 4m \pm 1$. We select $\{\{(i-1)(p-1)2p+\{X_1, X_2, \cdots, X_{\frac{p}{2}}, ((p-1)(2p-1)+1)-\{X_1, X_2, \cdots, X_{\frac{p-2}{2}}\}\}\}_{i=1}^{\frac{n}{2}}, \{(np(p-1)-ja)+\{X_1, X_2, \cdots, X_{\frac{a-1}{2}}\}\}_{j=0}^{\frac{p-2}{a}-1}, \{(np+1)(p-1)+1+4k(p-1)\}_{k=0}^{m-1}\}$ regions so that after making region crossing changes at these regions, the closure of the resultant braid is a trivial link by Theorem 4.1 and Lemma 3.1. Observe that the number of selected regions is $\frac{n}{2}(\sum\frac{p}{2}+\sum\frac{p-2}{2})+\frac{p-2}{a}\sum\frac{a-1}{2}+m=n\frac{p^2}{8}+\frac{p-2}{a}(\frac{a^2-1}{8})+\lfloor\frac{a+2}{4}\rfloor$. Thus,

$$u_R(K(p, np+a)) \leq n\frac{p^2}{8}+\frac{p-2}{a}\left(\frac{a^2-1}{8}\right)+\left\lfloor\frac{a+2}{4}\right\rfloor.$$

□

Theorem 4.4. *For torus links $K(p, np+a)$ where a is even, n is odd and $p \equiv 0 \pmod a$, we obtain the following:*

$$u_R(K(p, np+a)) \leq \frac{np^2+ap}{8}.$$

Proof. Consider $K(p, np+a)$ as the closure of $(\sigma_1\sigma_2\cdots\sigma_{p-1})^{np+a}$ as in Figure 1(a). Then, we select $\{\{(i-1)(p-1)2p+\{X_1, X_2, \cdots, X_{\frac{p}{2}}, ((p-1)(2p-1)+1)-\{X_1, X_2, \cdots, X_{\frac{p-2}{2}}\}\}\}_{i=1}^{\frac{n-1}{2}}, (n-1)p(p-1)+\{X_1, X_2, \cdots, X_{\frac{p}{2}}, \{ja+(p+a-1)(p-1)+1-\{X_1, X_2, \cdots, X_{\frac{a-2}{2}}\}\}_{j=1}^{\frac{p}{a}-1}\}\}$ regions so that after making region crossing changes at these regions, the closure of the resultant braid is a trivial link by Theorem 4.1 and Lemma 3.2. Observe that the number of regions selected is $\frac{n-1}{2}\left(\sum(\frac{p}{2})+\sum(\frac{p-2}{2})\right)+\sum(\frac{p}{2})+\frac{p}{a}\sum(\frac{a-2}{2}) = \frac{(n-1)p^2}{8}+\frac{p(p+2)}{8}+\frac{p}{a}\left(\frac{a(a-2)}{8}\right)$. Thus,

$$u_R(K(p, np+a)) \leq \frac{np^2+ap}{8}.$$ □

Theorem 4.5. *For torus links $K(p, np-1)$ with $n \geq 2$*

1. *If p is odd, then*
$$u_R(K(p, np-1)) \leq \frac{n(p^2-1)}{8}.$$
2. *If p is even and n is odd, then*
$$u_R(K(p, np-1)) \leq \frac{np^2-2p}{8}.$$
3. *If both p and n are even, then*
$$u_R(K(p, np-1)) \leq \frac{np^2}{8}.$$

Proof. Consider $K(p, np-1)$ as the closure of $(\sigma_1\sigma_2\cdots\sigma_{p-1})^{np-1}$ as in Figure 1(a). Then,
Case 1. If p is odd, then we select the regions $\{(i-1)(p-1)p+\{X_1, X_2, \cdots, X_{\frac{p-1}{2}}\}\}_{i=1}^{n-1}, ((n-1)p-1)(p-1)+\{X_1, X_2, \cdots, X_{\frac{p-1}{2}}\}$ so that after making region crossing change on these regions, we obtain

$$(\mu_1\mu_2\cdots\mu_p)^{n-1}\mu_2\mu_3\cdots\mu_p$$

where $\mu_i = \sigma_1\sigma_2\cdots\sigma_{p-i}\sigma_{p-i+1}^{-1}\sigma_{p-i+2}^{-1}\cdots\sigma_{p-1}^{-1}$. By Theorem 3.1 and Remark 1, the closure of $(\mu_1\mu_2\cdots\mu_p)^{n-1}\mu_2\mu_3\cdots\mu_p$ is a trivial knot.

Observe that number of crossing changes is $(n-1)\sum\frac{p-1}{2}+\sum\frac{p-1}{2} = \frac{n(p^2-1)}{8}$. Thus,

$$u_R(K(p, np+1)) \leq \frac{n(p^2-1)}{8}$$

Case 2. If p is even and n is odd, then we write $K(p, np-1) = K(p, (n-1)p+(p-1))$, where both p and $n-1$ are even. This result follows from Theorem 4.2.
Case 3. If both p and n are even, we select the regions same as we selected for $K(p, np)$ in Theorem 4.1. After making region crossing changes on selected regions, we are left with

$$(\mu_1\mu_2\cdots\mu_p\nu_1\nu_2\cdots\nu_p)^{\frac{n-2}{2}}\mu_1\mu_2\cdots\mu_p\nu_1\nu_2\cdots\nu_{p-1}.$$

By Theorem 3.1, Remark 1 and Lemma 3.1, its closure is unknot. Observe that the number of region crossing changes is $n\left(\frac{p^2}{8}\right)$. Thus,

$$u_R(K(p, np)) \leq \frac{np^2}{8}.$$ □

Theorem 4.6. *For torus links $K(p, np+2)$ with $n \geq 1$*

1. *If p is odd, then*
$$u_R(K(p, np+2)) \leq \frac{n(p^2-1)}{8} + \left\lfloor \frac{p+1}{4} \right\rfloor.$$
2. *If p is even and n is odd, then*
$$u_R(K(p, np+2)) \leq \frac{np^2+2p}{8}.$$
3. *If both p and n are even, then*
   (a) *if $p = 4m+2$ for some m, then $K(p, np+2)$ is not proper.*
   (b) *if $p = 4m$ for some m, then*
$$u_R(K(p, np+2)) \leq \frac{np^2+2p}{8}.$$

PROOF. Consider $K(p, np+2)$ as the closure of $(\sigma_1\sigma_2\cdots\sigma_{p-1})^{np+2}$ as in Figure 1(a). Then,
Case 1. If p is odd then $\exists\ m \in \mathbb{N} \cup \{0\}$ such that $p = 4m \pm 1$. We select $\{(i-1)(p-1)p + \{X_1, X_2, \cdots, X_{\frac{p-1}{2}}\}\}_{i=1}^{n}, \{(np+1)(p-1) - 4j\}_{j=0}^{m-1}$ regions so that after making region crossing change on these regions, we obtain by Theorem 3.1 and Lemma 3.3 that

$$\sigma_1\sigma_2\sigma_3\sigma_4^{-1}\sigma_1\sigma_2\sigma_3^{-1}\sigma_4^{-1};\ if\ p = 4m+1,$$
$$\sigma_1\sigma_2^{-1}\sigma_1^{-1}\sigma_2^{-1};\ if\ p = 4m-1.$$

We observe that the closure of these braids are trivial links.

Observe that the number of region crossing changes is $n\sum(\frac{p-1}{2}) + m = n(\frac{p^2-1}{8}) + \lfloor\frac{p+1}{4}\rfloor$. Thus,

$$u_R(K(p, np+2)) \leq \frac{n(p^2-1)}{8} + \left\lfloor \frac{p+1}{4} \right\rfloor.$$

Case 2. If p is even and n is odd, then by Theorem 4.4:

$$u_R(K(p, np+2)) \leq \frac{np^2+2p}{8}.$$

Case 3. If both p and n are even and $p \equiv 0\ (mod\ 4)$, then we select the regions $\{(i-1)(p-1)2p + \{X_1, X_2, \cdots, X_{\frac{p}{2}}, ((p-1)(2p-1)+1) - \{X_1, X_2, \cdots, X_{\frac{p-2}{2}}\}\}\}_{i=1}^{\frac{n}{2}}, \{(np+1)(p-1) - 4j\}_{j=0}^{\frac{p}{4}-1}$ so that by Theorem 3.1 and Lemma 3.3, after making region crossing change on these regions, we obtain

$$\sigma_1\sigma_2\sigma_3^{-1}\sigma_1\sigma_2^{-1}\sigma_3^{-1}.$$

We observe that the closure of this braid is a trivial link. Also, the number of region crossing changes is $\frac{n}{2}\left(\sum\frac{p}{2} + \sum\frac{p-2}{2}\right) + \frac{p}{4} = n(\frac{p^2}{8}) + \frac{p}{4}$. Thus,

$$u_R(K(p, np+2)) \leq \frac{np^2+2p}{8}.$$

But if $p \equiv 2\ (mod\ 4)$, then it follows from Theorem 2.1 that $K(p, np+2)$ is not proper. □

Theorem 4.7. *Consider torus links $K(p, np+a)$ where a is odd and $p \equiv -2\ (mod\ a)$, $n \geq 1$. Then we obtain the following:*

1. *If p is odd, then*

$$u_R(K(p, np+a)) \leq \frac{n(p^2-1)}{8} + \left(\frac{p+2}{a} - 1\right)\left(\frac{a^2-1}{8}\right) + \frac{(a-2)^2-1}{8} + \left\lfloor \frac{a}{4} \right\rfloor.$$

2. *If both p and n are even, then*

$$u_R(K(p, np+a)) \leq \frac{np^2}{8} + \left(\frac{p+2}{a} - 1\right)\left(\frac{a^2-1}{8}\right) + \frac{(a-2)^2-1}{8} + \left\lfloor \frac{a}{4} \right\rfloor.$$

PROOF. Consider $K(p, np+a)$ as the closure of $(\sigma_1\sigma_2\cdots\sigma_{p-1})^{np+a}$ as in Figure 1(a). Then,
Case 1. If p is odd, then we select $\{\{(i-1)(p-1)p + \{X_1, X_2, \cdots, X_{\frac{p-1}{2}}\}\}_{i=1}^{n}, \{(np(p-1) - ja) + \{X_1, X_2, \cdots, X_{\frac{a-1}{2}}\}\}_{j=0}^{\frac{p+2}{a}-2}, \{(np(p-1) - (\frac{p+2}{a} - 1)a) + \{X_1, X_2, \cdots, X_{\frac{a-3}{2}}\}\}, \{(np+a-2)(p-1) + (a-3) - 4l\}_{l=0}^{\lfloor \frac{a}{4} \rfloor}$ regions so that after making region crossing changes at these regions, the closure of the resultant braid is a trivial link by Theorem 4.1, Lemma 3.1 and Theorem 4.6. Observe that the number of selected regions is $n\sum(\frac{p-1}{2}) + \left(\frac{p+2}{a} - 1\right)\left(\sum(\frac{a-1}{2})\right) + \sum(\frac{a-3}{2}) + \lfloor\frac{a}{4}\rfloor = \frac{n(p^2-1)}{8} + \left(\frac{p+2}{a} - 1\right)\left(\frac{a^2-1}{8}\right) + \frac{(a-2)^2-1}{8} + \lfloor\frac{a}{4}\rfloor$. Thus,

$$u_R(K(p, np+a)) \leq \frac{n(p^2-1)}{8} + \left(\frac{p+2}{a} - 1\right)\left(\frac{a^2-1}{8}\right) + \frac{(a-2)^2-1}{8} + \left\lfloor \frac{a}{4} \right\rfloor.$$

Case 2. If both p and n are even, then we select $\{\{(i-1)(p-1)2p + \{X_1, X_2, \cdots, X_{\frac{p}{2}}, ((p-1)(2p-1)+1) - \{X_1, X_2, \cdots, X_{\frac{p-2}{2}}\}\}\}_{i=1}^{\frac{n}{2}}, \{(np(p-1) - ja) + \{X_1, X_2, \cdots, X_{\frac{a-1}{2}}\}\}_{j=0}^{\frac{p+2}{a}-2}, \{(np(p-1) - (\frac{p+2}{a} - 1)a) + \{X_1, X_2, \cdots, X_{\frac{a-3}{2}}\}\}, \{(np+a-2)(p-1) + (a-3) - 4l\}_{l=0}^{\lfloor \frac{a}{4} \rfloor}$ regions so that after making region crossing changes at these regions, the closure of the resultant braid is a trivial link by Theorem 4.1, Lemma 3.1 and Theorem 4.6. Observe that the number of selected regions is $\frac{n}{2}\left(\sum\frac{p}{2} + \sum\frac{p-2}{2}\right) + \left(\frac{p+2}{a} - 1\right)\left(\sum(\frac{a-1}{2})\right) + \sum(\frac{a-3}{2}) + \lfloor\frac{a}{4}\rfloor = \frac{np^2}{8} + \left(\frac{p+2}{a} - 1\right)\left(\frac{a^2-1}{8}\right) + \frac{(a-2)^2-1}{8} + \lfloor\frac{a}{4}\rfloor$. Thus,

$$u_R(K(p, np+a)) \leq \frac{np^2}{8} + \left(\frac{p+2}{a} - 1\right)\left(\frac{a^2-1}{8}\right) + \frac{(a-2)^2-1}{8} + \left\lfloor \frac{a}{4} \right\rfloor. \qquad \square$$

Theorem 4.8. *For torus links $K(p, np-2)$ with $n \geq 2$, we obtain the following*

1. *If p is odd, then*

$$u_R(K(p, np-2)) \leq \frac{n(p^2-1)}{8} - \frac{p-1}{2} + \left\lfloor \frac{p}{4} \right\rfloor.$$

2. *If p is even and n is odd, then*

$$u_R(K(p, np-2)) \leq \frac{np^2-2p}{8}.$$

3. *If both p and n are even, then*
    (a) *if $p \equiv 2$ (mod 4), then $K(p, np-2)$ is not proper.*
    (b) *if $p \equiv 0$ (mod 4), then*

$$u_R(K(p, np-2)) \leq \frac{np^2-2p}{8}.$$

Proof. Consider $K(p, np-2)$ as the closure of $(\sigma_1\sigma_2\cdots\sigma_{p-1})^{np-2}$ as in Figure 1(a). Then
Case 1. If p is odd, then result hold by Theorem 4.3.
Case 2. If p is even and n is odd, then select the regions $\{(i-1)(p-1)2p+\{X_1, X_2, \cdots, X_{\frac{p}{2}}, ((p-1)(2p-1)+1)-\{X_1, X_2, \cdots, X_{\frac{p-2}{2}}\}\}\}_{i=1}^{\frac{n-1}{2}}, ((n-1)p-1)(p-1)+\{X_1, X_2, \cdots, X_{\frac{p-2}{2}}\}$. Then after making region crossing change on these regions, we obtain

$$(\mu_1\mu_2\cdots\mu_p\nu_1\nu_2\cdots\nu_p)^{\frac{n-1}{2}}\mu_2\mu_3\cdots\mu_{p-1}$$

where $\mu_i = \sigma_1\sigma_2\cdots\sigma_{p-i}\sigma_{p-i+1}^{-1}\sigma_{p-i+2}^{-1}\cdots\sigma_{p-1}^{-1}$ and $\nu_i = \sigma_1^{-1}\sigma_2^{-1}\cdots\sigma_{p-i}^{-1}\sigma_{p-i+1}\cdots\sigma_{p-1}$. By Theorem 3.1, Remark 1 and Lemma 3.5, the closure of resultant braid is a trivial link.

Observe that the number of region crossing changes is equal to $\frac{(n-1)}{2}(\sum\frac{p}{2}+\sum\frac{p-2}{2})+\sum(\frac{p-2}{2}) = \frac{np^2-2p}{8}$. Thus,

$$u_R(K(p, np+1)) \leq \frac{np^2-2p}{8}.$$

Case 3. If both p and n are even, then

1. if $p = 4m+2$ for some m, then it is clear from Theorem 2.1 that $K(p, np-2)$ is not proper.
2. if $p = 4m$ for some m, then we select the regions $\{(i-1)(p-1)2p+\{X_1, X_2, \cdots, X_{\frac{p}{2}}, ((p-1)(2p-1)+1)-\{X_1, X_2, \cdots, X_{\frac{p-2}{2}}\}\}\}_{i=1}^{\frac{n-2}{2}}, (n-2)p(p-1)+\{X_1, X_2, \cdots, X_{\frac{p}{2}}, (2p-3)(p-1)+2-\{X_1, X_2, \cdots, X_{\frac{p-4}{2}}\}, \{(p+2+4i)(p-1)+1\}_{i=0}^{m-2}\}$. After making region crossing changes on selected regions, we are left with

   $$(\mu_1\mu_2\cdots\mu_p\nu_1\nu_2\cdots\nu_p)^{\frac{n-2}{2}}\mu_1\mu_2\cdots\mu_p\beta$$

   where $\beta = \beta_1\beta_2\cdots\beta_{p-2}$ and β_i is defined as: $\beta_1 = \sigma_1^{-1}\sigma_2^{-1}\cdots\sigma_{p-1}^{-1}$, $\beta_{p-2} = \sigma_1\sigma_2\cdots\sigma_{p-1}$ and for the remaining $i^{'s}$,
   if $i \equiv 1\ (mod\ 4)$, $\beta_i = \sigma_1\sigma_2^{-1}\sigma_3^{-1}\sigma_4^{-1}\cdots\sigma_{p-i}^{-1}\sigma_{p-i+1}\cdots\sigma_{p-1}$,
   if $i \equiv 2\ (mod\ 4)$, $\beta_i = \sigma_1\sigma_2\sigma_3^{-1}\sigma_4^{-1}\cdots\sigma_{p-1}^{-1}\sigma_{p-i+1}\cdots\sigma_{p-1}$ and
   if $i \equiv 0$ *or* $3\ (mod\ 4)$, $\beta_i = \sigma_1^{-1}\sigma_2\sigma_3^{-1}\sigma_4^{-1}\cdots\sigma_{p-i}^{-1}\sigma_{p-i+1}\cdots\sigma_{p-1}$.
   By Theorem 3.1, Remark 1 and Lemma 3.2, the closure of

   $$(\mu_1\mu_2\cdots\mu_p\nu_1\nu_2\cdots\nu_p)^{\frac{n-2}{2}}\mu_1\mu_2\cdots\mu_p\beta \sim \underbrace{\sigma_1^{-1}}_{1}\underbrace{\sigma_1}_{2}\underbrace{\sigma_1^{-1}}_{3}\underbrace{\sigma_1^{-1}}_{4}\underbrace{\sigma_1}_{5}\cdots\underbrace{\sigma_1^{-1}}_{p-2}$$

   is a trivial link.
   Observe that the number of region crossing changes is $\frac{n-2}{2}\left(\sum\frac{p}{2}+\sum\frac{p-2}{2}\right)+\sum(\frac{p}{2})+\sum(\frac{p-4}{2})+\frac{p}{4}-1 = \frac{np^2-2p}{8}$. Thus,

   $$u_R(K(p, np)) \leq \frac{np^2-2p}{8}. \qquad \square$$

Theorem 4.9. *For torus links $K(p, np+3)$ with $n \geq 1$*

1. *If p is odd, then*

   $$u_R(K(p, np+3)) \leq \frac{n(p^2-1)}{8}+\left\lfloor\frac{p+1}{3}\right\rfloor.$$

2. *If p is even and n is odd, then*

$$u_R(K(p, np+3)) \leq \frac{np^2+2p}{8} + \left\lfloor \frac{p+2}{6} \right\rfloor.$$

3. *If both p and n are even, then*

$$u_R(K(p, np+3)) \leq \frac{np^2}{8} + \left\lfloor \frac{p+1}{3} \right\rfloor.$$

Proof. Consider $K(p, np+3)$ as the closure of $(\sigma_1\sigma_2\cdots\sigma_{p-1})^{np+3}$ as in Figure 1(a). Then
Case 1. If p is odd, then the result follows from Theorem 4.2.
Case 2. If p is even and n is odd, then $p+2 = 6m+a$ for some $m \geq 0$ and $0 \leq a < 6$ and we select the regions $\{(i-1)(p-1)2p+\{X_1, X_2, \cdots, X_{\frac{p}{2}}, ((p-1)(2p-1)+1)-\{X_1, X_2, \cdots, X_{\frac{p-2}{2}}\}\}\}_{i=1}^{\frac{n-1}{2}}, (n-1)p(p-1)+\{X_1, X_2, \cdots, X_{\frac{p}{2}}\}, \{(np+2)(p-1)-2-6i\}_{i=0}^{m-1}$. Then after making region crossing change on these regions, we obtain

$$(\mu_1\mu_2\cdots\mu_p\nu_1\nu_2\cdots\nu_p)^{\frac{n-1}{2}}\mu_1\mu_2\cdots\mu_{p-1}\mu_p\beta,$$

where $\mu_i = \sigma_1\sigma_2\cdots\sigma_{p-i}\sigma_{p-i+1}^{-1}\sigma_{p-i+2}^{-1}\cdots\sigma_{p-1}^{-1}$, $\nu_i = \sigma_1^{-1}\sigma_2^{-1}\cdots\sigma_{p-i}^{-1}\sigma_{p-i+1}\cdots\sigma_{p-1}$ and $\beta = \sigma_1^{-1}\sigma_2^{-1}\cdots\sigma_{p-1}^{-1}\sigma_1^{g_{2,1}}\sigma_2^{g_{2,2}}\cdots\sigma_{p-1}^{g_{2,p-1}}\sigma_1^{g_{3,1}}\sigma_2^{g_{3,2}}\cdots\sigma_{p-1}^{g_{3,p-1}}$ with $g_{2,j} = -1$ for $j = \{\{(p-2),(p-3)\}-6i\}_{i=0}^{m-1}$ and $g_{3,j} = -1$ for $j = \{\{(p-3),(p-4)\}-6i\}_{i=0}^{m-1}$.
By Theorem 3.1 and Lemma 3.4, the closure of the resultant braid is a trivial link.

Observe that the number of region crossing changes is equal to $\frac{(n-1)}{2}(\sum\frac{p}{2}+\sum\frac{p-2}{2})+\sum(\frac{p}{2})+m = \frac{np^2+2p}{8}+\lfloor\frac{p+2}{6}\rfloor$. Thus,

$$u_R(K(p, np+3)) \leq \frac{n(p^2+2p)}{8} + \left\lfloor \frac{p+2}{6} \right\rfloor.$$

Case 3. If both p and n are even, then the result follows from Theorem 4.2. ☐

Lemma 4.1. *The closure of the braid*
$\underbrace{\sigma_i^{-1}\sigma_{i-1}^{-1}\sigma_{i-2}^{-1}\sigma_{i-3}^{-1}}_{1}\underbrace{\sigma_{i+1}^{-1}\sigma_i^{-1}\sigma_{i-1}^{-1}\sigma_{i-2}^{-1}}_{2}\underbrace{\sigma_{i+2}\sigma_{i+1}\sigma_i\sigma_{i-1}}_{3}\underbrace{\sigma_{i+3}^{-1}\sigma_{i+2}^{-1}\sigma_{i+1}\sigma_i}_{4}\underbrace{\sigma_{i+4}^{-1}\sigma_{i+3}^{-1}\sigma_{i+2}\sigma_{i+1}}_{5}\underbrace{\sigma_{i+5}\sigma_{i+4}\sigma_{i+3}\sigma_{i+2}}_{6}$
$\underbrace{\sigma_{i+6}^{-1}\sigma_{i+5}^{-1}\sigma_{i+4}\sigma_{i+3}}_{7}\underbrace{\sigma_{i+7}\sigma_{i+6}\sigma_{i+5}\sigma_{i+4}}_{8}$ *is trivial link.*

Proof. The proof follows elimentary braid relations.

Theorem 4.10. *For torus links $K(p, np+4)$ with $n \geq 1$*

1. *If p is odd, then*
   (a) *if $p \equiv 1$ or 3 (mod 8), then*

$$u_R(K(p, np+4)) \leq \frac{n(p^2-1)}{8} + \left\lfloor \frac{p}{2} \right\rfloor.$$

   (b) *if $p \equiv 5$ or 7 (mod 8), then*

$$u_R(K(p, np+4)) \leq \frac{n(p^2-1)}{8} + \left\lceil \frac{p}{2} \right\rceil.$$

2. *If both* p *and* n *are even, then*
    (a) *if* $p \equiv 0, 2$ *or* 6 $(mod\ 8)$*, then*
    $$u_R(K(p, np+4)) \leq \frac{np^2}{8} + \frac{p}{2}.$$
    (b) *if* $p \equiv 4$ $(mod\ 8)$*, then* $K(p, np)$ *is not proper.*
3. *If* p *is even and* n *is odd, then*
    (a) *if* $p \equiv 0$ $(mod\ 4)$*, then*
    $$u_R(K(p, np+4)) \leq \frac{np^2}{8} + \frac{p}{2}.$$
    (b) *if* $p \equiv 2$ $(mod\ 4)$*, then* $K(p, np+4)$ *is not proper.*

Proof. Consider $K(p, np{+}4)$ as the closure of $(\sigma_1\sigma_2\cdots\sigma_{p-1})^{np}(\sigma_1\sigma_2\sigma_3)^4\sigma_4\sigma_3\cdots\sigma_1\sigma_5\sigma_4\cdots\sigma_2\cdots\sigma_{p-1}$ $\sigma_{p-2}\cdots\sigma_{p-4}$ as in Figure 1(b). Also, if $p \equiv a$ $(mod\ 8)$, then $\exists\ m \in \mathbb{N} \cup \{0\}$ such that $p = mp + a$. Then

Case 1. If p is odd, then

1. if $p \equiv 1$ $(mod\ 8)$, then we select $\{\{(i-1)(p-1)p + \{X_1, X_2, \cdots, X_{\frac{p-1}{2}}\}\}_{i=1}^{n}, np(p-1) + \{6, 10, 12, 18, \{24i + \{1, 3, 12, 18\}\}_{i=1}^{m-1}\}\}$ regions.
2. if $p \equiv 3$ $(mod\ 8)$, then we select $\{\{(i-1)(p-1)p + \{X_1, X_2, \cdots, X_{\frac{p-1}{2}}\}\}_{i=1}^{n}, np(p-1) + \{6, 10, 12, 18, \{24i + \{1, 3, 12, 18\}\}_{i=1}^{m-1}, 24m + \{3\}\}\}$ regions.
3. if $p \equiv 5$ $(mod\ 8)$ and $m \neq 0$, then we select $\{\{(i-1)(p-1)p + \{X_1, X_2, \cdots, X_{\frac{p-1}{2}}\}\}_{i=1}^{n}, np(p-1) + \{6, 10, 12, 18, \{24i + \{1, 3, 12, 18\}\}_{i=1}^{m-1}, 24m + \{1, 3, 12\}\}\}$ regions and if $m = 0$, then we select $\{\{(i-1)(p-1)p + \{X_1, X_2, \cdots, X_{\frac{p-1}{2}}\}\}_{i=1}^{n}, np(p-1) + \{6, 10, 12\}\}$ regions.
4. if $p \equiv 7$ $(mod\ 8)$ and $m \neq 0$, then we select $\{\{(i-1)(p-1)p + \{X_1, X_2, \cdots, X_{\frac{p-1}{2}}\}\}_{i=1}^{n}, np(p-1) + \{6, 10, 12, 18, \{24i + \{1, 3, 12, 18\}\}_{i=1}^{m}\}\}$ regions and if $m = 0$, then we select $\{\{(i-1)(p-1)p + \{X_1, X_2, \cdots, X_{\frac{p-1}{2}}\}\}_{i=1}^{n}, np(p-1) + \{6, 10, 12, 18\}\}$ regions.

Case 2. If both p and n are even, then

1. if $p \equiv 0$ $(mod\ 8)$, then we select $\{\{(i-1)(p-1)2p + \{X_1, X_2, \cdots, X_{\frac{p}{2}}, ((p-1)(2p-1)+1) - \{X_1, X_2, \cdots, X_{\frac{p-2}{2}}\}\}\}_{i=1}^{\frac{n}{2}}, np(p-1) + \{6, 10, 12, 18, \{24i + \{1, 3, 12, 18\}\}_{i=1}^{m-1}\}\}$ regions.
2. if $p \equiv 2$ $(mod\ 8)$, then we select $\{\{(i-1)(p-1)2p + \{X_1, X_2, \cdots, X_{\frac{p}{2}}, ((p-1)(2p-1)+1) - \{X_1, X_2, \cdots, X_{\frac{p-2}{2}}\}\}\}_{i=1}^{\frac{n}{2}}, np(p-1) + \{6, 10, 12, 18, \{24i + \{1, 3, 12, 18\}\}_{i=1}^{m-1}, 24m + \{3\}\}\}$ regions.
3. if $p \equiv 6$ $(mod\ 8)$ and $m \neq 0$, then we select $\{\{(i-1)(p-1)2p + \{X_1, X_2, \cdots, X_{\frac{p}{2}}, ((p-1)(2p-1)+1) - \{X_1, X_2, \cdots, X_{\frac{p-2}{2}}\}\}\}_{i=1}^{\frac{n}{2}}, np(p-1) + \{6, 10, 12, 18, \{24i + \{1, 3, 12, 18\}\}_{i=1}^{m-1}, 24m + \{3\}\}\}$ regions and if $m = 0$, then we select $\{\{(i-1)(p-1)2p + \{X_1, X_2, \cdots, X_{\frac{p}{2}}, ((p-1)(2p-1)+1) - \{X_1, X_2, \cdots, X_{\frac{p-2}{2}}\}\}\}_{i=1}^{\frac{n}{2}}, np(p-1) + \{6, 10, 12\}\}$ regions.
4. if $p \equiv 4$ $(mod\ 8)$, then it is not proper follows from Theorem 2.1.

In all the above cases, after making region crossing changes the closure of the resultant braid is trivial link by Theorem 4.1 and Lemma 4.1.

Case 3. If p is even and n is odd, then

1. if $p \equiv 0$ $(mod\ 4)$, then the result follows from Theorem 4.4.
2. if $p \equiv 2$ $(mod)4$, then the result follows from Theorem 2.1. □

Theorem 4.11. *For torus links $K(p, np+5)$ with $n \geq 1$*

1. *If p is odd, and*
   (a) *if $p \equiv 0$ or ± 1 $(mod\ 5)$, then*
   $$u_R(K(p, np+5)) \leq \frac{n(p^2-1)}{8} + 3\left\lfloor \frac{p+1}{5} \right\rfloor.$$
   (b) *if $p \equiv 2$ $(mod\ 5)$, then*
   $$u_R(K(p, np+5)) \leq \frac{n(p^2-1)}{8} + \frac{3p-1}{5}.$$
   (c) *if $p \equiv 3$ $(mod\ 5)$, then*
   $$u_R(K(p, np+5)) \leq \frac{n(p^2-1)}{8} + \frac{3p+1}{5}.$$
2. *If both p and n are even, and*
   (a) *if $p \equiv 0$ or ± 1 $(mod\ 5)$, then*
   $$u_R(K(p, np+5)) \leq \frac{np^2}{8} + 3\left\lfloor \frac{p+1}{5} \right\rfloor.$$
   (b) *if $p \equiv 2$ $(mod\ 5)$, then*
   $$u_R(K(p, np+5)) \leq \frac{np^2}{8} + \frac{3p-1}{5}.$$
   (c) *if $p \equiv 3$ $(mod\ 5)$, then*
   $$u_R(K(p, np+5)) \leq \frac{np^2}{8} + \frac{3p+1}{5}.$$

PROOF. The proof directly follows from Theorem 4.2, Theorem 4.3 and Theorem 4.7. □

5. Conclusion

It would be interesting to extend the results obtained here to find exact region unknotting number for torus knots/links. This seems to be a hard problem. In [2], Ayaka showed that $u_R(K) \leq c(K)/2 + 1$, where $C(K)$ is the number of crossings of the knot K. Here, the sharp bounds provided for torus knots/links are much smaller than the upper bound $c(K)/2+1$. In particular, whatever be the case we considered in this paper, the upper bound we obtain, for $u_R(K)$, lies between $c/8$ and $c/5$ and as $p \to \infty$, the sharp upper bound provided tends to $c/8$. In case of proper $K(p, np)$ and $K(p, np+1)$, we conjecture that the sharp upper bound provided in the paper is equal to the region unknotting number for them.

Acknowledgements

The first author thanks CSIR New Delhi and IIT Ropar for providing financial assistance and research facilities.

References

[1] V. Siwach and P. Madeti, A method for unknotting torus knots. math.GT/1207.4918v1, 2012.

[2] Ayaka Shimizu, Region crossing change is an unknotting operation. math.GT/1011.6304v2, 2010.

[3] Ayaka Shimizu, On region unknotting numbers. RIMS Kokyuroku 1766, 15-22, 2011.

[4] H. Murakami, Some metrics on classical knots. Math. Ann. 270, 35-45, 1985.

[5] P. Kronheimer and T. Mrowka, Gauge theory for embedded surfaces, I, Topology, 32 , 773-826. 1993.

[6] P. Kronheimer and T. Mrowka, Gauge theory for embedded surfaces, II, Topology, 34 , 37-97, 1995.

[7] H. Murakami, Y. Nakanishi, On a certain move generating link-homology. Math. Ann. 284, 75-89, 1989.

[8] Cheng Zhiyun, Gao Hongzhu, On region crossing change and incidence matrix, Science China Mathematics. Vol. 55 No. 7. 14871495, 2012.

[9] Y. Nakanishi, Replacements in the Conway third identity, Tokyo J. Math. 14, 197-203, 1991.

[10] Haruko Aida, Unknotting operation for Polygonal type, Tokyo J. Math. Vol. 15, No. 1, 111-121, 1992.

[11] J. Hoste, Y. Nakanishi and K. Taniyama, Unknotting operations involving trivial tangles, Osaka J. Math. 27 , 555-566, 1990.

[12] Cheng Zhiyun, When is region crossing change an unknotting operation? math.GT/1201.1735v1, 2012.